Multiparametric imaging of adhesive nanodomains at the surface of *Candida albicans* by atomic force microscopy

Cécile Formosa-Dague, Marion L Schiavone, Anita Boisrame, Mathias L Richard, Raphael E Duval, Etienne Dague

► To cite this version:

Cécile Formosa-Dague, Marion L Schiavone, Anita Boisrame, Mathias L Richard, Raphael E Duval, et al.. Multiparametric imaging of adhesive nanodomains at the surface of *Candida albicans* by atomic force microscopy. *Nanomedicine: Nanotechnology, Biology and Medicine*, 2015, 11 (1), pp.57-65. 10.1016/j.nano.2014.07.008 . hal-01553144

HAL Id: hal-01553144

<https://hal.laas.fr/hal-01553144>

Submitted on 7 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multiparametric imaging of adhesive nanodomains at the surface of *Candida albicans* by Atomic Force Microscopy

Cécile Formosa, MS^{1,2,3,4}, Marion Schiavone, MS^{1,2}, Anita Boisrame, PhD^{5,6}, Mathias L. Richard, PhD^{5,6}, Raphaël E. Duval, PhD^{3,4,7}, and Etienne Dague, PhD^{1,2}

¹ CNRS, LAAS, 7 avenue du Colonel Roche, F-31400 Toulouse, France

² Université de Toulouse ; LAAS, F31400 Toulouse, France

³ CNRS, UMR 7565, SRSMC, Vandœuvre-lès-Nancy, France

⁴ Université de Lorraine, UMR 7565, Faculté de Pharmacie, Nancy, France

⁵ INRA, UMR1319 Micalis, F-78352 Jouy-en-Josas, France

⁶ AgroParisTech, UMR Micalis, F-78850 Thiverval Grignon, France

⁷ ABC Platform®, Nancy, France

Corresponding author: Etienne Dague, LAAS-CNRS, 7 av du Colonel Roche, 31400 Toulouse, France. Phone number: +33 5 61 33 78 41 Mail : edague@laas.fr

Abstract word count: 138 words / Manuscript word count: 4451 words / Number of figures: 5 / Number of references: 40

This work was supported by an ANR young scientist program (AFMYST project ANR-11-JSV5-001-01 n° SD 30024331) to ED. CF and MS are respectively supported by a grant from “Direction Générale de l’Armement” (DGA) and from Lallemand SAS.

Abstract

Candida albicans is an opportunistic pathogen. It adheres to mammalian cells through a variety of adhesins that interact with hosts ligands. The spatial organization of these adhesins on the cellular interface is however poorly understood, mainly because of the lack of an instrument able to track single molecules on single cells. In this context, the Atomic Force Microscope (AFM) makes it possible to analyze the force signature of single proteins on single cells. The present study is dedicated to the mapping of the adhesive properties of *C. albicans* cells. We observed that the adhesins at the cell surface were organized in nanodomains composed of free or aggregated mannoproteins. This was demonstrated by the use of functionalized AFM tips and synthetic amyloid forming/disrupting peptides. This direct visualization of amyloids nanodomains will help in understanding the virulence factors of *C. albicans*.

Background

The yeast *Candida albicans* has emerged as a major public health problem these last two decades. This opportunistic pathogen causes a wide range of infections from surface infections, to mucosal and blood-stream infections¹. Whereas mucosal infections are common and occur in healthy organisms, blood-stream infections are observed only in immunocompromised patients and are life-threatening. This type of infections, also known as candidaemia, can develop into disseminated candidiasis when the infection spreads to internal organs, leading to high mortality rates². In order to colonize and subsequently to disseminate in the blood stream *C. albicans* needs to adhere to different substrates. This first stage of infection³ is mediated by adhesins that are found on the surface of the yeast cell wall. Many of these adhesins are mannoproteins, and among them, the adhesin family identified as having a major role in host cell attachment is the Als (Agglutinin-like Sequences) family⁴.

The Als were initially reported as having homologies with the proteins responsible for auto-agglutination in the baker yeast *Saccharomyces cerevisiae*. Eight Als have been identified, they all are primarily involved in host-pathogen interactions⁵. It was found that there were amyloid-forming sequences in the Als adhesins of *Candida albicans*⁶. Amyloids are insoluble fibrillar protein aggregates whose core consists in crystalline arrays of identical sequence in many molecules of the amyloid protein^{7,8}. Cells expressing the Als proteins can rapidly aggregate, and the aggregation has amyloid-like properties. Like amyloid formation, aggregation ability propagates through the adherent cell population and depends on conformational changes of the Als protein. This transition of the conformational state to an aggregative state of the proteins is characterized by the formation of hydrophobic nanodomains on the entire surface of the cell⁹.

A few papers written by Lipke's team were dedicated to the direct visualization of these nanodomains using fluorescent dyes such as thioflavin T or 8-anilino-1-naphthalene-sulfonic acid

(ANS)^{6,8,9}. Another technique that can be used to visualize these nanodomains is Atomic Force Microscopy (AFM). AFM has recently emerged as a valuable tool to study the surface of living cells¹⁰, and especially pathogenic cells¹¹. This technology has been used by Alsteens *et al.* to image the formation and propagation of nanodomains in living yeast cells¹² and also to unfold amyloid proteins from the yeasts surface using Single Molecule Force Spectroscopy^{13–15}. To this end, the authors functionalized AFM tips with antibodies targeted against the Als protein directly or against an epitope tag present in the Als protein. These studies allowed the authors to localize the adhesive nanodomains caused by the aggregation of Als proteins at the surface of living yeast cells, and to unravel the structure of the Als proteins studied by stretching.

In our study, we used AFM as an imaging tool to visualize and localize adhesins nanodomains at the surface of living wild-type *Candida albicans* cells. Using recent developments in the AFM technology, we have imaged and quantified at the same time the nanomechanical properties, the adhesiveness (force and nature of the interaction), the size and the thickness of the nanodomains^{16,17}, at high resolution. The data collected showed that these nanodomains are localized differently at the surface of the cell, depending on the structures featured by the cells (bud scars, buds). We also showed that there were degrees of adhesiveness, depending on whether the amyloid proteins had totally aggregated (hydrophobic nanodomains) or not, and that these degrees of aggregation were directly correlated to the stiffness of the yeast cell wall. Finally, using force measurements and amyloid forming or inhibiting peptides, we showed that Als proteins (probably among others) were participating to these nanodomains.

Methods

Yeasts growth conditions

Candida albicans (from ABC Platform® Bugs Bank, Nancy, France) was stocked at -80°C, revived on Yeast Peptone Dextrose agar (Difco, 242720-500g) and grown in Yeast Peptone Dextrose broth (Difco, 242820-500g) for 20 hours at 30°C under static conditions.

Sample preparation for AFM experiments

Yeast cells were concentrated by centrifugation, washed two times in acetate buffer (18 mM CH₃COONa, 1 mM CaCl₂, 1 mM MnCl₂, pH = 5.2), resuspended in acetate buffer, and immobilized on polydimethylsiloxane (PDMS) stamps prepared as described by Dague *et al*¹⁸. Briefly, freshly oxygen activated microstructured PDMS stamps were covered by a total of 100 µL of the solution of cells and allowed to stand for 15 minutes at room temperature. The cells were then deposited into the microstructures of the stamp by convective/capillary assembly. Images were recorded in acetate buffer in Quantitative Imaging™ mode with MLCT AUWH (Bruker) cantilevers (nominal spring constant of 0.01 N/m). The applied force was kept at 1.5 nN for imaging and at 0.5 nN for force spectroscopy experiments. The loading rate for imaging was of 2 500 000 pN/s (acquisition frequency of the force curves is of 25 Hz) and for force spectroscopy of 75 000 pN/s (acquisition frequency of the force curves is of 1.25 Hz). For imaging and force spectroscopy, we used an AFM Nanowizard III (JPK Instruments, Berlin, Germany). The cantilevers spring constants were determined by the thermal noise method¹⁹. For all the results presented in this study, silicon nitride AFM tips were bare, except in the case of figure 4g (lower panel), where a functionalized AFM tip has been used.

AFM tips fonctionnalization

The functionalized tips were produced according to a french patent of the authors described later in sensors and actuators²⁰. Briefly, AFM tips were functionalized with dendrimers presenting

CHO functions able to covalently link with NH₂ functions of proteins. These dendritips were then incubated with the lectin Concanavalin A (Sigma, L7647-100MG, 100µg/mL) for 1 hour, before being used for force spectroscopy experiments.

Results analysis

All results were analyzed using the Data Processing software from JPK Instruments. The stiffness values measured on cells were determined from the slope of the linear portion of the raw deflections versus piezo displacement curves, according to:

$$k_{cell} = k \left(\frac{s}{1-s} \right)$$

with s the experimentally accessible slope of the compliance region reached for sufficient loading forces. In this model, the experimental setup can be represented by two linear springs, one is the AFM's cantilever, and the other is the cell envelope exhibiting an effective spring constant. It is then possible to calculate the effective spring constant k_{cell} of the cell envelope from the observed slope s of the force curve and the known spring constant k of the cantilever²¹.

Results

Candida albicans cells display localized adhesiveness

Thanks to our innovative method to immobilize cells into PDMS stamps¹⁸, and using Quantitative ImagingTM mode¹⁶, we were able to image and quantify the adhesive properties of single *C. albicans* cells at the same time. Figure 1a shows a budding yeast cell; on the corresponding adhesion image (Figure 1b), we can see that only the bud, and not the mother-cell, presents adhesives patches. This original result is surprising as non-budding cells are highly adhesive (see below). This result seems to indicate that the mother-cell cell wall changes during

the budding process. As for the cell in figure 1c, this cell displays two bud scars, a common feature at the surface of yeast cells, which are not adhesive whereas the rest of the cell is. This type of distribution of the adhesion on yeast cells has already been seen using immunofluorescence with antibodies targeted against surface proteins of *C. albicans*. Coleman *et al.* for example showed that the Als1 protein was expressed all over *C. albicans* cells, with the exception of bud scars²². The comparison of our results to these data suggests then that the adhesions probed by AFM might be due to surface proteins, such as Als1 in the case of the cell presenting bud scars, but perhaps also others adhesins.

***C. albicans* cell wall adhesins are able to aggregate into nanodomains**

As showed before^{9,12}, the proteins expressed at the surface of *C. albicans* cell wall are able to aggregate, and to form nanodomains. However, these nanodomains have not been yet characterized at the nanoscale, nor were imaged at high resolution. In fact, these nanodomains have specific adhesive properties that can be mapped using AFM in the Quantitative ImagingTM mode. High resolution (256 pixels²) adhesive images are presented in Figure 2. It shows adhesive nanodomains, at the surface of a living wild-type *C. albicans* cell. On the cell presented in this figure (Figure 2a), the corresponding adhesion image shows very distinct adhesive nanodomains that were probed with bare AFM tips. These nanodomains are homogeneously distributed all over the cell here, which does not present any morphological features such as buds or bud scars. When zooming into small areas on top of the cell (white squares on figure 2b), we could measure the area of each nanodomain. On this cell and on another one showed in Figure 3a, 60 nanodomains areas were measured; the values obtained plotted on figure 2e shows that nanodomains have an average area of $0.09 \pm 0.03 \mu\text{m}^2$. This corresponds to an average diameter of 170 nm, which confirms the nanoscale of these nanodomains. Some of the nanodomains are also higher than the

rest of the cell wall. When the whole cell is imaged, it is not visible; however, specific analysis of the Figure 3d and the graphic 3h representing the topography of the cell surface revealed nanodomains that had a different height compared to the rest of the cell. The cross-sections taken along the blue line showed the height of a nanodomain of 20 nm. Once again, this confirmed the nanoscale of the nanodomains at the surface of *C. albicans*.

Different nanodomains have different nanomechanical properties

Adhesion is measured as the rupture force recorded when retracting the tip from the surface, when approaching and pulling with the tip on the cell wall, thus AFM makes it possible to measure nanomechanical properties of living cells. Here we choose to use an analysis based on the Hooke model which considers the coupled cantilever / cell wall as a spring. The stiffness values measured on cells were determined from the slope of the linear portion of the raw deflections versus piezo displacement curves, according to:

$$k_{cell} = k \left(\frac{s}{1-s} \right)$$

with s the experimentally accessible slope of the compliance region reached for sufficient loading forces. Indeed, the most interesting result in this study is the correlation that can be directly made between the adhesiveness of the nanodomain, and its stiffness. Nanodomains on figure 3e (adhesion map) circled in red were found on the stiffness image (figure 3f) circled in black; they correspond to the zones where the stiffness of the cell wall is increased, to 13.4 ± 0.3 nN/ μ m. As for less adhesive nanodomains, they do not present any difference in stiffness from the rest of the cell, and are 12.4 ± 0.2 nN/ μ m. The 3D-view of the adhesion, mapped with the stiffness (figure 3g) illustrates this clear correlation; the more adhesive the nanodomain is, the stiffer it is.

Another fascinating point is that for the more adhesive nanodomains, the retract force curves present typical hydrophobic adhesions^{23–25}, with adhesions occurring immediately when the tip is retracted from the surface. Force curves from the other, less adhesive, nanodomains presented retract adhesions resembling to proteins unfolding, occurring several nanometers after the tip withdrawal. Therefore it seems that the nanodomains are of 2 different natures. There is a class of nanodomains, hydrophobic, higher and stiff, and another class displaying proteins unfolding properties, as soft as the rest of the cell wall. What is the molecular nature of these 2 types of nanodomains, and are they correlated?

Understanding the adhesive properties of the 2 nanodomains classes

To answer the previous question, we monitored the retract force curves recorded on the nanodomains (Figure 4). We found the same correlation as in figure 3; the force curves recorded on an adhesive nanodomains presented hydrophobic retract adhesions, whereas the force curves recorded on a less adhesive nanodomain presented protein, unfolding like, profiles. In order to determine the nature of these last unfoldings, we probed the surface of *C. albicans* cells with an AFM tip functionalized with Concanavalin A (ConA), a protein that interacts with yeast mannoproteins, such as surface adhesins. The resulting force curves (figure 4g) showed retract adhesions displaying unfoldings of different lengths, but with a similar profile. We also observed condensed spikes with adhesion forces between 0 and 50 pN. This value was consistent with specific interactions between ConA and mannoproteins²⁶.

In a previous study conducted in 2009¹⁴ by Alsteens *et al.*, adhesins (Als5) were unfolded from the surface of live *S. cerevisiae* cells overexpressing this protein. The retract force curves obtained in this study show high similarity with the ones we obtain here with functionalized AFM tips, with the presence of serin-threonin rich segments (condensed pikes on figure 4g). We can

therefore, based on this comparison with the data of the literature, conclude that the less adhesive nanodomains at the surface of live *C. albicans* are composed of free adhesins, and maybe of Als proteins. However, since all adhesins (like Als, Hwp1, Eap1, Rbt1 etc) are mannoproteins, we cannot, at this stage, make a statement on which adhesins are unfolded here.

As for the second type of nanodomains, the hydrophobic ones, our hypothesis is that they are composed of the same proteins as the less adhesive ones. In fact, adhesins (like Als) display amyloids sequences located on a domain of the protein called T, that enable them to change their conformation⁶ and to aggregate into amyloid nanodomains. And when this phenomena is started, it propagates to the whole cell⁹. We therefore made the hypothesis that the adhesive nanodomains are in fact amyloid nanodomains, made of Als proteins.

From adhesins to amyloid nanodomains: the role of Als proteins

To verify this hypothesis, and according to the literature on Als proteins, we synthesized a peptide exhibiting the same sequence as the one of the T domain of the Als1/3/5 proteins. We then put this peptide in the presence of the cells in order to trigger the amyloids formation. We also synthesized the same peptide, but with a mutation on one amino acid (V326N peptide), in order to obtain a peptide that inhibits the formation of the amyloid nanodomains⁸. Since Als3 is only expressed on the surface of hyphae, we will only be able to generate or destroy the amyloid formation of Als 1 and 5. The results presented in figure 5 showed a cell before and after adding the mutated peptide. We can clearly see on these adhesion images the loss of general adhesion and of two nanodomains at the center of the cell. It seems like the mutated peptide disrupted the amyloids at the surface of the cell. And the other way around, when cells were incubated with the amyloid forming peptide, we could observe the formation of the nanodomains at the surface of the cells, as it is showed on the adhesion images on local areas on top of *C. albicans* cells in

figure 5c, d and e. These results allow us to conclude that the proteins at the origin of the nanodomains are then mannoproteins, more specifically Als1 or 5, or both, that form amyloids.

Discussion

We show in this study that wild-type live *C. albicans* cells exhibit extraordinary adhesive properties. In the case of budding cells, placed in acetate buffer at 25°C for 2h, we observed that the mother cell is not adhesive and that only the bud presents adhesive nanodomains. On the contrary, we show that non budding cells are covered by adhesive nanodomains, in the same experimental conditions. This illustrates the amazing plasticity of this species^{27,28} able to grow as a commensal or as a pathogen^{29,30}, in all the parts of the intestinal track, but also on the vaginal mucosa, as unicellular budding cells or as filamentous hyphae. Moreover its cell wall is permanently remodeled as a reaction to its environment (temperature, pH, dissolved O₂, ions, interacting surface/cells/bacteria) what makes it challenging to reproduce the experimental conditions inducing a certain cell wall phenotype. We then demonstrate that the molecules at the origin of these adhesions could aggregate into nanodomains, which can be probed at high resolution using a suited AFM mode, QITM. These nanodomains are different in terms of level of adhesiveness, which is a property directly correlated to their stiffness and to the hydrophobic state or not of the molecule at the origin of these nanodomains. We then went further in the study, using functionalized AFM tips, and were able to determine that the less adhesive nanodomains were formed by mannoproteins that can interact specifically with Concanavalin A. These mannoproteins are able to aggregate to form the adhesive nanodomains because they have amyloid properties as we showed in figure5.

Amyloid aggregation is a primitive³¹ and very stable³² protein folding and a common structural motif. It is a cross β -sheet quaternary structure that usually auto-aggregates as fibrils. It

has been, first, associated with neurodegenerative diseases like Alzheimer, Parkinson, or Creutzfeldt-Jakob diseases. However it is more and more unclear if the amyloids lesions are the cause or a consequence of the disease. Amyloid aggregates are now described as functional proteins assembly and can be found from bacteria to humans³³. In microorganisms, amyloid has been described as a functional coat³⁴. It consists in curli (*E. coli*), chaperons (*Streptomyces*) or hydrophobins (*Aspergillus etc*); all of these proteins are implicated in adhesion to the host and in the invasion, infection process. It is now well known that adhesins (and especially *Als*) of *C. albicans* have amyloid-forming sequences^{6,35} and that these proteins form domains involved in cell aggregation or biofilm formation³⁶. Nevertheless the characterization, structure and properties of the amyloids adhesive nanodomains remain unclear.

In this work we measured for the first time the nanoscale size of amyloid domains (average area of $0.09 \mu\text{m}^2$) at the surface of live *C. albicans* cells. The domains are of 2 different classes. Some present the characteristic of individual proteins whereas the others are hydrophobic, stiffer than the rest of the cell ($13.4 \pm 0.3 \text{ nN}/\mu\text{m}$ compared to $12.4 \pm 0.2 \text{ nN}/\mu\text{m}$), and are slightly protruding. It means that there is a state modification from soluble proteins into insoluble proteins, which is a characteristic of amyloid structures. This transformation is dependent on the proteins concentration and can only occur when the protein density exceeds a threshold. The roles of the two classes of domains are probably different. On one hand we could hypothesize that the hydrophobic nanodomains were involved in the cell adhesion to abiotic hydrophobic surfaces or to cell membrane as it is known that membrane binding is an inherent property of amyloid aggregates^{37,38}. Amyloid aggregation is also a way to store proteins, in a limited space and to sort them when required. This has been demonstrated for hormones in secretory granules^{39,40}. Thus *C. albicans* may store some adhesins for the subsequent invasion phases. On the other hand the protein like domains may be responsible for specific adhesion to

fibronectin and other extra cellular proteins of the matrix. It seems rational that several adhesins, brought together, would be more efficient in a binding process than a single adhesion. This finding has to be added to *C. albicans* plasticity²⁸ and participate to explain its remarkable adaptation and pathogenicity.

However, there are still many things to explore on the cell wall of *C. albicans*, and future work will be dedicated to exploring the changes appearing on the mother cell during the budding process.

References

1. Gow NA, Hube B. Importance of the *Candida albicans* cell wall during commensalism and infection. *Curr Opin Microbiol.* 2012;**15**:406–412.
2. Poulain D. *Candida albicans* , plasticity and pathogenesis. *Crit Rev Microbiol.* 2013:1–10.
3. Naglik JR, Moyes DL, Wächtler B, Hube B. *Candida albicans* interactions with epithelial cells and mucosal immunity. *Microbes Infect.* 2011;**13**:963–976.
4. Hoyer LL. The ALS gene family of *Candida albicans*. *Trends Microbiol.* 2001;**9**:176–180.
5. Hoyer LL, Green CB, Oh S-H, Zhao X. Discovering the secrets of the *Candida albicans* agglutinin-like sequence (ALS) gene family – a sticky pursuit. *Med Mycol.* 2008;**46**:1–15.
6. Ramsook CB, Tan C, Garcia MC, Fung R, Soybelman G, Henry R, et al. Yeast Cell Adhesion Molecules Have Functional Amyloid-Forming Sequences. *Eukaryot Cell.* 2010;**9**:393–404.
7. Sawaya MR, Sambashivan S, Nelson R, Ivanova MI, Sievers SA, Apostol MI, et al. Atomic structures of amyloid cross- β spines reveal varied steric zippers. *Nature.* 2007;**447**:453–457.
8. Garcia MC, Lee JT, Ramsook CB, Alsteens D, Dufrêne YF, Lipke PN. A Role for Amyloid in Cell Aggregation and Biofilm Formation. *PLoS ONE.* 2011;**6**:e17632.
9. Rauceo JM, Gaur NK, Lee K-G, Edwards JE, Klotz SA, Lipke PN. Global Cell Surface Conformational Shift Mediated by a *Candida albicans* Adhesin. *Infect Immun.* 2004;**72**:4948–4955.

- 300 10. Müller DJ, Dufrêne YF. Atomic force microscopy: a nanoscopic window on the cell
301 surface. *Trends Cell Biol.* 2011;**21**:461–469.
- 302 11. Alsteens D, Beaussart A, El-Kirat-Chatel S, Sullan RMA, Dufrêne YF. Atomic Force
303 Microscopy: A New Look at Pathogens. *PLoS Pathog.* 2013;**9**:e1003516.
- 304 12. Alsteens D, Garcia MC, Lipke PN, Dufrêne YF. Force-induced formation and propagation
305 of adhesion nanodomains in living fungal cells. *Proc Natl Acad Sci.* 2010;**107**:20744 –
306 20749.
- 307 13. Beaussart A, Alsteens D, El-Kirat-Chatel S, Lipke PN, Kucharíková S, Van Dijck P, et al.
308 Single-Molecule Imaging and Functional Analysis of Als Adhesins and Mannans during
309 *Candida albicans* Morphogenesis. *ACS Nano.* 2012;**6**:10950–10964.
- 310 14. Alsteens D, Dupres V, Klotz SA, Gaur NK, Lipke PN, Dufrêne YF. Unfolding Individual
311 Als5p Adhesion Proteins on Live Cells. *ACS Nano.* 2009;**3**:1677–1682.
- 312 15. Alsteens D, Ramsook CB, Lipke PN, Dufrêne YF. Unzipping a Functional Microbial
313 Amyloid. *ACS Nano.* 2012;**6**:7703–7711.
- 314 16. Chopinet L, Formosa C, Rols MP, Duval RE, Dague E. Imaging living cells surface and
315 quantifying its properties at high resolution using AFM in QITM mode. *Micron.* 2013;**48**:26–
316 33.
- 317 17. Dufrêne YF, Martínez-Martín D, Medalsy I, Alsteens D, Müller DJ. Multiparametric
318 imaging of biological systems by force-distance curve-based AFM. *Nat Methods.*
319 2013;**10**:847–854.
- 320 18. Dague E, Jauvert E, Laplatine L, Viallet B, Thibault C, Ressler L. Assembly of live micro-
321 organisms on microstructured PDMS stamps by convective/capillary deposition for AFM
322 bio-experiments. *Nanotechnology.* 2011;**22**.
- 323 19. Hutter JL, Bechhoefer J. Calibration of atomic-force microscope tips. *Rev Sci Instrum.*
324 1993;**64**:1868–1873.
- 325 20. Jauvert E, Dague E, Séverac M, Ressler L, Caminade A-M, Majoral J-P, et al. Probing
326 single molecule interactions by AFM using bio-functionalized dendritips. *Sens Actuators B*
327 *Chem.* 2012;**168**:436–441.
- 328 21. Arnoldi M, Fritz M, Bäuerlein E, Radmacher M, Sackmann M, Boulbitch A. Bacterial
329 turgor pressure can be leasured by atomic force microscopy. *Phys Rev E.* 1999;**62**:1034-1044
330
- 331 22. Coleman DA, Oh S-H, Zhao X, Hoyer LL. Heterogeneous distribution of *Candida albicans*
332 cell-surface antigens demonstrated with an Als1-specific monoclonal antibody.
333 *Microbiology.* 2010;**156**:3645–3659.
- 334 23. Dague E, Alsteens D, Latge JP, Verbelen C, Raze D, Baulard AR, et al. Chemical force
335 microscopy of single live cells. *Nano Lett.* 2007;**7**:3026–3030.

- 336 24. Alsteens D, Dague E, Rouxhet PG, Baulard AR, Dufrene YF. Direct measurement of
337 hydrophobic forces on cell surfaces using AFM. *Langmuir*. 2007;**23**:11977–11979.
- 338 25. Dague E, Alsteens D, Latgé J-P, Dufrêne YF. High-Resolution Cell Surface Dynamics of
339 Germinating *Aspergillus fumigatus* Conidia. *Biophys J*. 2008;**94**:656–660.
- 340 26. Alsteens D, Dupres V, Evoy KM, Wildling L, Gruber HJ, Dufrêne YF. Structure, cell wall
341 elasticity and polysaccharide properties of living yeast cells, as probed by AFM.
342 *Nanotechnology*. 2008;**19**.
- 343 27. Odds F. *Candida* and candidosis. A review and bibliography. 2nd edition. London: Bailliere
344 Tindall; 1988.
- 345 28. Poulain D. *Candida albicans* , plasticity and pathogenesis. *Crit Rev Microbiol*. 2013:1–10.
- 346 29. Gow NA, Hube B. Importance of the *Candida albicans* cell wall during commensalism and
347 infection. *Curr Opin Microbiol*. 2012;**15**:406–412.
- 348 30. Gow NAR, van de Veerdonk FL, Brown AJP, Netea MG. *Candida albicans* morphogenesis
349 and host defence: discriminating invasion from colonization. *Nat Rev Micro*. 2012;**10**:112–
350 122.
- 351 31. Greenwald J, Riek R. Biology of Amyloid: Structure, Function, and Regulation. *Structure*.
352 2010;**18**:1244–1260.
- 353 32. Perczel A, Hudáky P, Pálfi VK. Dead-End Street of Protein Folding: Thermodynamic
354 Rationale of Amyloid Fibril Formation. *J Am Chem Soc*.; **129**:14959–14965.
- 355 33. Fowler DM, Koulov AV, Balch WE, Kelly JW. Functional amyloid--from bacteria to
356 humans. *Trends Biochem Sci*. 2007;**32**:217–224.
- 357 34. Gebbink MFBG, Claessen D, Bouma B, Dijkhuizen L, Wösten HAB. Amyloids — a
358 functional coat for microorganisms. *Nat Rev Microbiol*. 2005;**3**:333–341.
- 359 35. Otoo HN, Lee KG, Qiu W, Lipke PN. *Candida albicans* Als Adhesins Have Conserved
360 Amyloid-Forming Sequences. *Eukaryot Cell*. 2008;**7**:776–782.
- 361 36. Garcia MC, Lee JT, Ramsook CB, Alsteens D, Dufrêne YF, Lipke PN. A Role for Amyloid
362 in Cell Aggregation and Biofilm Formation. *PLoS ONE*. 2011;**6**:e17632.
- 363 37. Sparr E, Engel MFM, Sakharov DV, Sprong M, Jacobs J, de Kruijff B, et al. Islet amyloid
364 polypeptide-induced membrane leakage involves uptake of lipids by forming amyloid
365 fibers. *FEBS Lett*. 2004;**577**:117–120.
- 366 38. Gellermann GP, Appel TR, Tannert A, Radestock A, Hortschansky P, Schroeckh V, et al.
367 Raft lipids as common components of human extracellular amyloid fibrils. *Proc Natl Acad*
368 *Sci U S A*. 2005;**102**:6297–6302.

- 369 39. Maji SK, Perrin MH, Sawaya MR, Jessberger S, Vadodaria K, Rissman RA, et al.
370 Functional Amyloids As Natural Storage of Peptide Hormones in Pituitary Secretory
371 Granules. *Science*. 2009;**325**:328–332.
- 372 40. Dannies PS. Concentrating hormones into secretory granules: layers of control. *Mol Cell*
373 *Endocrinol*. 2001;**177**:87–93.

374

375 **Figure Caption**

376 **Figure 1. Localization of the adhesive properties of *C. albicans* cells.** (a) Height image (z-
377 range = 1.5 μm) of a budding *C. albicans* cell in a PDMS stamp, and (b) adhesion image
378 corresponding to the height image. On (a), MC stands for Mother cell, BC stands for Budding
379 cell, and the red dotted line represents the demarcation between the two different cells. (c) Height
380 image (z-range = 3.5 μm) of a single *C. albicans* cell exhibiting two bud scars, and (d) adhesion
381 image corresponding to the height image.

382

383 **Figure 2. Imaging of the adhesive domains of *C. albicans* cells in acetate buffer, at 25°C for**
384 **2 hours.** (a) Height image (z-range = 2.5 μm) of a single *C. albicans* cell in a
385 polydimethylsiloxane (PDMS) stamp, and (b) adhesion images corresponding to the height
386 images. (c, d) Adhesion images of small areas on top of the cell, represented by the white squares
387 in b. (e) distribution of the areas values of the domains in c and d.

388

389 **Figure 3. Nanomechanics of the adhesive domains of *C. albicans* cells.** (a) Height image (z-
390 range = 2.5 μm) of a *C. albicans* cell in a PDMS stamp, (b) corresponding adhesion image, and
391 (c) corresponding stiffness image. (d) Height image (z-range = 100 nm) of a small area on top of
392 the cell, represented by the white square on (a), (e) corresponding adhesion image and (f)
393 corresponding stiffness image. Note that the adhesive nanodomains circled in red on (e) are also

found on the stiffness image (black circles on f). (g) is a 3D-image of the adhesion mapped with the stiffness. (h) cross-section taken along the blue line on (d), and (i), distribution of the stiffness values corresponding to the yeast cell wall and the less adhesive domains (blue columns) or to the most adhesive domains (yellow columns).

Figure 4. Adhesion force curves of *C. albicans* adhesive domains. (a) Height image (z-range = 4.0 μm) of a *C. albicans* cell in a PDMS stamp. (b) Adhesion image of a small area on top of the cell, represented by the white square on (a). (c and d) representative force curves obtained on the zones indicated by the arrows on (c). (e) Height image (z-range = 2.5 μm) of a *C. albicans* cell in a PDMS stamp and (f) corresponding adhesion image recorded with a bare tip. (g) representative force curves obtained in the zone delimited by a white square on (f) with the Con A tip.

Figure 5. Imaging of the adhesive domains of *C. albicans* cells treated with Als1, 3, 5p amyloid disrupting peptide (V326N peptide) or Als1, 3, 5p amyloid forming peptide (Als peptide). (a) Adhesion image of a single *C. albicans* cell in a PDMS stamp and (b) corresponding adhesion image after adding V326N peptide. (c, d and e) Adhesion images of small areas on top of a *C. albicans* cell after adding the Als peptide.