

Stability analysis of a 1D wave equation with a nonmonotone distributed damping

Swann Marx, Yacine Chitour, Christophe Prieur

► **To cite this version:**

Swann Marx, Yacine Chitour, Christophe Prieur. Stability analysis of a 1D wave equation with a nonmonotone distributed damping. 2019. <hal-02006292>

HAL Id: hal-02006292

<https://hal.laas.fr/hal-02006292>

Submitted on 5 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stability analysis of a 1D wave equation with a nonmonotone distributed damping

Swann Marx ^{*,****,†}, Yacine Chitour ^{**}, Christophe Prieur ^{***}

^{*} LAAS-CNRS, Université de Toulouse, CNRS, 7 avenue du colonel Roche, 31400, Toulouse, France marx.swann@gmail.com.

^{**} Laboratoire des Signaux et Systèmes (L2S), CNRS - CentraleSupélec - Université Paris-Sud, 3, rue Joliot Curie, 91192, Gif-sur-Yvette, France, yacine.chitour@lss.supelec.fr.

^{***} Univ. Grenoble Alpes, CNRS, Grenoble INP, Gipsa-lab, F-38000 Grenoble, France, christophe.prieur@gipsa-lab.fr

^{****} DeustoTech, University of Deusto, 48007 Bilbao, Basque Country, Spain.

[†] Facultad de Ingeniería, Universidad de Deusto, Avda. Universidades, 24, 48007, Bilbao - Basque Country - Spain.

Abstract: This paper is concerned with the asymptotic stability analysis of a one dimensional wave equation subject to a nonmonotone distributed damping. A well-posedness result is provided together with a precise characterization of the asymptotic behavior of the trajectories of the system under consideration. The well-posedness is proved in the nonstandard L^p functional spaces, with $p \in [2, \infty]$, and relies mostly on some results collected in Haraux (2009). The asymptotic behavior analysis is based on an attractivity result on a specific infinite-dimensional linear time-variant system.

Keywords: 1D wave equation, nonlinear control, Lyapunov functionals.

1. INTRODUCTION

This paper is concerned with the asymptotic behavior of a one-dimensional wave equation subject to a nonmonotone nonlinear damping. For control systems, it is always crucial to consider nonlinear feedback laws. In particular, the definition of the nonlinearity under consideration in this paper includes the saturation, which models amplitude limitations on the actuator. Such a phenomenon appears in most of control systems, and it can lead to undesirable behavior in terms of stability. Moreover, considering that such a nonlinearity is nonmonotone is crucial for control systems, because the model of the nonlinearity might have some error.

However, as illustrated in many papers (Alabau-Boussouira (2012), Haraux (2009), Marx et al. (2018), Slemrod (1989), Feireisl (1993), etc.), the monotone property of the nonlinearity is crucial to first prove the asymptotic stability of the system under consideration and second characterize the trajectory of the latter system. In order to characterize the asymptotic behavior of the trajectory of the system under consideration, we consider the initial conditions in an another functional setting than the classical one, that is $L^p(0, 1)$, with $p \in [2, \infty]$.

There exists a vast litterature about linear PDEs subject to monotone nonlinear dampings. For instance, in Slemrod (1989), asymptotic stability of the origin of abstract con-

trol systems subject to monotone nonlinear dampings is proved, using an infinite-dimensional version of LaSalle's invariance principle. These results have been then extended to more general infinite-dimensional systems in Marx et al. (2017a). More recently, in Marx et al. (2018), trajectory of such systems have been characterized via Lyapunov techniques. Let us also mention Prieur et al. (2016) and Marx et al. (2017b), where a wave equation and a nonlinear Korteweg-de Vries, respectively, subject to a nonlinear monotone damping are considered and where the global asymptotic stability is proved. Finally, in Kang and Fridman (2017), the local asymptotic stability of a heat equation coupled with an ODE, controlled from the boundary with a saturated feedback law, is proved.

The case of nonmonotone nonlinear damping have been considered in few papers. In Feireisl (1993), the global asymptotic stability of a one-dimensional wave equation subject to a nonmonotone nonlinear damping is proved, thanks to some compensated compactness technique. The characterization of the trajectories is, however, not provided. In Martinez and Vancostenoble (2000), the trajectory of a wave equation in two dimensions subject to a nonmonotone damping is characterized, but for only a specific nonmonotone damping. In this paper, we rather focus on a more general nonlinear nonmonotone damping, but only for a one dimensional wave equation. We are able to characterize the trajectories of the system by studying the functional spaces $L^p(0, 1)$.

There exists also few papers dealing with the one dimensional wave equation in this functional setting. Let us

* This works has been partially supported by Advanced Grant DYCON (Dynamic Control) of the European Research Council Executive Agency.

mention Haraux (2009), which derives a well-posedness analysis of a one dimensional wave equation in this functional setting. Moreover, an optimal decay rate is obtained for this equation. Recently, in Amadori et al. (2019), a similar result have been obtained, using techniques coming from conservation laws theory. Note that both of these results hold true only for monotone nonlinear damping, which is not the case of our paper.

In our paper, after introducing a general nonlinear non-monotone damping, we propose a well-posedness analysis of a one-dimensional wave equation subject to such a damping. Futhermore, using Lyapunov techniques, we are able to characterize the trajectory of this system in the L^p , $p \in [2, \infty)$, provided that the initial conditions are in L^∞ . This proof is mainly based on a result about linear time-varying infinite-dimensional system, which is introduced and proved in Appendix A.

This paper is organized as follows. In Section 2, the main results of the paper are collected. We propose a well-posedness and an asymptotic stability theorems. The second result proposes futhermore a precise characterization of the trajectories of the system. Section 3 is devoted to the proof of the main results. Finally, Section 4 collects some concluding remarks, together with further research lines to be followed. Appendix A introduces a result of independant interest about exponential convergence of a specific time-varying infinite-dimensional linear systems, following a finite-dimensional strategy provided in Chitour et al. (1995) and Liu et al. (1996), but which is instrumental for the proof of our asymptotic stability theorems.

Notation: For any $p \in [2, \infty)$, the space $L^p(0, 1)$ denotes the space of functions f satisfying $\left(\int_0^1 |f(x)|^p dx\right)^{\frac{1}{p}} < +\infty$. The space $L^\infty(0, 1)$ denotes the space of functions satisfying $\text{ess sup}_{x \in [0,1]} |f(x)| \leq +\infty$. For any $p \in [2, \infty]$, the Sobolev space $W^{1,p}(0, 1)$ (resp. $W^{2,p}(0, 1)$) is defined as follows $W_0^{1,p}(0, 1) := \{f \in L^p(0, 1) \mid f' \in L^p(0, 1) \text{ and } f(0) = f(1) = 0\}$ (resp. $W^{2,p}(0, 1) := \{f \in L^p(0, 1) \mid f', f'' \in L^p(0, 1)\}$).

2. MAIN RESULTS

The aim of this paper is to provide an asymptotic stability analysis of the following system:

$$\begin{cases} z_{tt}(t, x) = z_{xx}(t, x) - \sigma(a(x)z_t(t, x)), & (t, x) \in \mathbb{R}_+ \times [0, 1] \\ z(t, 0) = z(t, 1) = 0, & t \in \mathbb{R}_+ \\ z(0, x) = z_0(x), z_t(0, x) = z_1(x), & x \in [0, 1], \end{cases} \quad (1)$$

where z denotes the state, $a : [0, 1] \rightarrow \mathbb{R}_+$ is measurable and bounded by some positive constant a_∞ and σ is a scalar nonlinear damping which satisfies the following properties:

Definition 1. [Scalar nonlinear damping] A function $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ is said to be a scalar damping function if

1. It is locally Lipschitz and odd;
2. One has $\sigma(0) = 0$;
3. For any $s \in \mathbb{R}$, $\sigma(s)s > 0$;
4. The function σ is differentiable at $s = 0$ with $\sigma'(0) = C_1$ for some $C_1 > 0$.

Due to this definition (especially, item 2), the origin is an equilibrium point for (1). Note that this nonlinearity is not assumed to be *monotone*. The monotone property is in many cases really useful for either the well-posedness of the equation or the asymptotic stability of the origin. We refer the reader to Marx et al. (2018) for a complete discussion on this topic.

Example 1. (Example of nonlinear dampings). Below are listed some examples of nonlinear dampings:

1. The classical saturation, defined as follows:

$$\sigma(s) = \text{sat}(s) := \begin{cases} s & \text{if } |s| \leq 1, \\ \frac{s}{|s|} & \text{if } |s| \geq 1, \end{cases} \quad (2)$$

satisfies all the properties of Definition 1.

2. The following nonlinearity

$$\sigma(s) = \text{sat}\left(\frac{1}{4}s - \frac{1}{30}\sin(10s)\right) \quad (3)$$

is also a nonlinear damping. Note moreover that it is not monotone, as illustrated by Figure 1.




Fig. 1. For any $s \in [-5, 5]$, the figure illustrates the function σ given by (3)

As illustrated in Marx et al. (2018), some regularity is needed to obtain a characterization of the asymptotic stability of (1). To be more precise, we need the state z_t to be bounded in $L^\infty(0, 1)$. With a monotone nonlinearity σ , one would have this regularity result thanks to some nonlinear semigroup theorems. In the case of the system under consideration in this paper, we need to follow another strategy.

Our strategy relies on the introduction of these functional spaces:

$$\begin{aligned} H_p(0, 1) &:= W_0^{1,p}(0, 1) \times L^p(0, 1) \\ D_p(0, 1) &:= W^{2,p}(0, 1) \cap W_0^{1,p}(0, 1) \times W_0^{1,p}(0, 1), \end{aligned} \quad (4)$$

with $p \in [2, \infty]$. The first functional space is equipped with the following norm:

$$\begin{aligned} \|(z, z_t)\|_{H_p(0,1)} &:= \left(\int_0^1 |z_x(x)|^p dx \right)^{\frac{1}{p}} \\ &\quad + \left(\int_0^1 |z_t(x)|^p dx \right)^{\frac{1}{p}}, \quad \forall p \in [2, \infty), \\ \|(z, z_t)\|_{H_\infty(0,1)} &:= \|z_x\|_{L^\infty(0,1)} + \|z_t\|_{L^\infty(0,1)}, \quad \text{for } p = \infty. \end{aligned} \quad (5)$$

The second one is equipped with the following norm

$$\|(z, z_t)\|_{D_p(0,1)} := \left(\int_0^1 |z_{xx}(x)|^p dx \right)^{\frac{1}{p}} + \left(\int_0^1 |z_{tx}(x)|^p dx \right)^{\frac{1}{p}}, \quad \forall p \in [2, \infty)$$

$$\|(z, z_t)\|_{D_\infty(0,1)} := \|z_{xx}\|_{L^\infty(0,1)} + \|z_{tx}\|_{L^\infty(0,1)}, \quad \text{for } p = \infty. \quad (6)$$

Considering that $(z_0, z_1) \in H_\infty(0, 1)$, our aim is to prove an L^∞ regularity on the state z_t so that we can obtain a characterization of the asymptotic behavior of the trajectory in $H_p(0, 1)$, with $p \in [2, \infty)$. But before stating such a result, we need a suitable notion of solution for systems modeled with (1). Such a notion is provided by the following well-posedness theorem:

Theorem 1. [Well-posedness] *For any initial conditions $(z_0, z_1) \in H_\infty(0, 1)$, there exists a unique solution $(z, z_t) \in L^\infty(\mathbb{R}_+; W^{1,\infty}(0, 1)) \times W^{1,\infty}(\mathbb{R}_+; L^\infty(0, 1))$ to (1). Moreover, the following inequality is satisfied, for all $t \geq 0$*

$$\|(z, z_t)\|_{H_\infty(0,1)} \leq 2 \max(\|z'_0\|_{L^\infty(0,1)}, \|z_1\|_{L^\infty(0,1)}). \quad (7)$$

Now that the functional setting is introduced, we are in position to state our asymptotic stability result

Theorem 2. [Semi-global exponential stability] *Consider initial conditions $(z_0, z_1) \in H_\infty(0, 1)$ satisfying:*

$$\|(z_0, z_1)\|_{H_\infty(0,1)} \leq R, \quad (8)$$

where R is a positive constant. Then, for any $p \in [2, \infty)$, there exist two positive constants $K := K(R)$ and $\beta := \beta(R)$ such that

$$\|(z, z_t)\|_{H_p(0,1)} \leq K e^{-\beta t} \|(z_0, z_1)\|_{H_p(0,1)}, \quad \forall t \geq 0. \quad (9)$$

3. PROOF OF THE MAIN RESULTS

3.1 Proof of Theorem 1

The proof of Theorem 1 relies on the following wave equation with a source term

$$\begin{cases} z_{tt}(t, x) = z_{xx}(t, x) + h(t, x), & (t, x) \in \mathbb{R}_+ \times [0, 1] \\ z(t, 0) = z(t, 1) = 0, & t \in \mathbb{R}_+ \\ z(0, x) = z_0(x), \quad z_t(0, x) = z_1(x), & x \in [0, 1], \end{cases} \quad (10)$$

where h denotes the source term. From (Haraux, 2018, Theorem 1.3.8), we know that, provided that $z_0, z_1 \in H_2$ and that $h \in L^2(\mathbb{R}_+; L^2(0, 1))$, there exists a unique solution $z \in C(\mathbb{R}_+; H_0^1(0, 1)) \cap C^1(\mathbb{R}_+; L^2(0, 1))$ to (10). In particular, since $H_\infty(0, 1) \subset H_2$, this result holds true also for initial conditions $(z_0, z_1) \in H_\infty$. The first step of our analysis in this section is to prove that picking initial conditions in $H_\infty(0, 1)$ and the source term $h \in L^2(\mathbb{R}_+; L^\infty(0, 1))$ improves also the regularity of the solution z itself.

To do so, our aim is to give an explicit formula for the latter equation, using the reflection method surveyed in Strauss (1992). Roughly speaking, this method consists in extending the explicit formulation of trajectory of the wave equation in an unbounded domain to a bounded domain. To do so, we extend the initial datas to the whole line to be odd with respect to both $x = 0$ and $x = 1$, that is

$$\tilde{z}_0(-x) = -\tilde{z}_0(x) \text{ and } \tilde{z}_0(2-x) = -\tilde{z}_0(x), \quad (11)$$

¹ The term z'_0 denotes the derivative of z_0 with respect to x .

where \tilde{z}_0 denotes the 2-periodic odd extension of z_0 . A way to do this is to define \tilde{z}_0 as follows:

$$\tilde{z}_0(x) = \begin{cases} z_0(x), & 0 < x < 1 \\ -z_0(-x), & -1 < x < 0 \\ \text{extended to be of period 2} \end{cases} \quad (12)$$

We can define similarly a 2-periodic odd extension of z_1 (resp. h), denoted by \tilde{z}_1 (resp. \tilde{h}). Thanks to (Strauss, 1992, Theorem 1, Page 69), we can therefore define the explicit trajectory z of (10) (known as the D'Alembert formula) as follows:

$$\begin{aligned} z(t, x) &= \frac{1}{2} [\tilde{z}_0(x+t) + \tilde{z}_0(x-t)] + \frac{1}{2} \int_{x-ct}^{x+ct} \tilde{z}_1(s) ds \\ &+ \frac{1}{2} \int_0^t \int_{x-(t-s)}^{x+(t-s)} \tilde{h}(w, s) dw ds. \end{aligned} \quad (13)$$

We can further define z_t as follows

$$\begin{aligned} z_t(t, x) &= \frac{1}{2} (\tilde{z}'_0(x+t) - \tilde{z}'_0(x-t)) \\ &+ \frac{1}{2} (\tilde{z}_1(x+t) - \tilde{z}_1(x-t)) \\ &+ \frac{1}{2} \int_0^t (\tilde{h}(s, x+(t-s)) - \tilde{h}(s, x-(t-s))) ds \end{aligned} \quad (14)$$

It is clear from these two latter equations that, when picking $(z_0, z_1) \in H_\infty(0, 1)$ and $h \in L^2(\mathbb{R}_+; L^\infty(0, 1))$, then

$$z \in C(\mathbb{R}_+; W^{1,\infty}(0, 1)) \cap C^1(\mathbb{R}_+; L^\infty(0, 1)).$$

We assume now that \tilde{h} is written as follows

$$\tilde{h}(t, x) := -\sigma(\tilde{a}(x)y(t, x)), \quad (15)$$

where \tilde{a} is a 2-periodic extension of a , $y \in L^2(0, T; L^2(0, 1))$ and σ is a scalar nonlinear damping (which is odd, due to Item 1 of Definition 1). In particular, it means that $h \in L^2(\mathbb{R}_+; L^2(0, 1))$. The proof of Theorem 1 consists first in applying a fixed-point theorem, which will allow us to prove the well-posedness of (1) for a small time $T > 0$ and second in using a stability result in Haraux (2009), stated as follows

Proposition 1. *Let us consider initial condition $z_0, z_1 \in H_\infty(0, 1)$. If there exists a solution to (1), then the time derivative of the following functional along the trajectories of (1)*

$$\phi(z, z_t) := \int_0^1 (F(z_t + z_x) + F(z_t - z_x)) dx, \quad (16)$$

with F any even and convex function, satisfies

$$\frac{d}{dt} \phi(z, z_t) \leq 0.$$

The latter proposition implies that

$$\phi(z, z_t) \leq \phi(z_0, z_1), \quad \forall t \geq 0. \quad (17)$$

In particular, following the discussion in the proof of Corollary 2.3. in Haraux (2009), if one picks ² $F(s) = [\text{Pos}(|s| - 2 \max(\|z_0\|_{L^\infty(0,1)}, \|z_1\|_{L^\infty(0,1)}))]^2$, then one obtains that

$$\phi(z, z_t) = 0, \quad (18)$$

² The function $\text{Pos} : s \in \mathbb{R} \rightarrow \text{Pos}(s) \in \mathbb{R}_+$ is defined as follows:

$$\text{Pos}(s) = \begin{cases} s & \text{if } s > 0 \\ 0 & \text{if } s \leq 0. \end{cases}$$

which implies that, for all $t \geq 0$

$$\max(\|z_x(t, \cdot)\|_{L^\infty(0,1)}, \|z_t(t, \cdot)\|_{L^\infty(0,1)}) \leq 2 \max(\|z'_0\|_{L^\infty(0,1)}, \|z_1\|_{L^\infty(0,1)}). \quad (19)$$

Noticing that

$$\|(z, z_t)\|_{H^\infty(0,1)} \leq \max(\|z_x(t, \cdot)\|_{L^\infty(0,1)}, \|z_t(t, \cdot)\|_{L^\infty(0,1)}),$$

it is clear then that, for all $t \geq 0$

$$\|(z, z_t)\|_{H^\infty(0,1)} \leq 2 \max(\|z'_0\|_{L^\infty(0,1)}, \|z_1\|_{L^\infty(0,1)}) \quad (20)$$

This estimate implies that, once one is able to prove that there exists a solution (z, z_t) of (1) in

$$L^\infty([0, T]; W^{1,\infty}(0, 1)) \cap W^{1,\infty}([0, T]; L^\infty(0, 1)),$$

for a small time $T > 0$, then the well-posedness of (1) is ensured in $L^\infty(\mathbb{R}_+; W^{1,\infty}(0, 1)) \cap W^{1,\infty}(\mathbb{R}_+; L^\infty(0, 1))$.

Now, we are in position to prove Theorem 1.

Proof. Let us define \mathcal{F}_T the space of measurable functions defined on $[0, T] \times \mathbb{R}$ which are bounded, odd and 2-periodic in space. We endow \mathcal{F}_T with the L^∞ -norm so that it becomes a Banach space. Hence, denoting by $\|\cdot\|_T$ the norm of the latter functional space, we have, for every $y \in \mathcal{F}_T$

$$\|y\|_T := \sup_{(t,x) \in [0,T] \times \mathbb{R}} |y(t,x)|. \quad (21)$$

Let us consider $\mathbf{B}_K(y)$ the closed ball in \mathcal{F}_T centered at $y \in \mathcal{F}_T$ of radius $K \geq 0$, where K remains to be defined.

We can now define the mapping with which we will apply a fixed-point

$$\begin{aligned} \phi_T : \mathcal{F}_T &\rightarrow \mathcal{F}_T \\ y &\mapsto \phi_T(y), \end{aligned} \quad (22)$$

where

$$\begin{aligned} \phi_T(y) &= \frac{1}{2} (\tilde{z}'_0(x+t) - \tilde{z}'_0(x-t)) + \frac{1}{2} (\tilde{z}_1(x+t) - \tilde{z}_1(x-t)) \\ &- \frac{1}{2} \int_0^t \left(\sqrt{\tilde{a}(x+t-s)} \sigma(\sqrt{\tilde{a}(x+t-s)}) y(s, x-(t-s)) \right. \\ &\left. + \sqrt{\tilde{a}(-x+t-s)} \sigma(\sqrt{\tilde{a}(-x+t-s)}) y(s, -x-(t-s)) \right) ds \end{aligned} \quad (23)$$

Using the fact that σ is locally Lipschitz, note that, for every $y_0 \in \mathcal{F}_T$ and $K > 0$, there exists a positive constant $C(y_0, K)$ such that, for every $y, \tilde{y} \in \mathbf{B}_K(y_0)$

$$\|\phi_T(y) - \phi_T(\tilde{y})\|_T \leq C(y_0, K)T \|y - \tilde{y}\|_T. \quad (24)$$

Pick K such that

$$K := 2(\|\phi_T(y_0)\|_T + \|y_0\|_T) \quad (25)$$

and T sufficiently small such that

$$C(y_0, K)T \leq 1. \quad (26)$$

Consider a sequence $(y_n)_{n \in \mathbb{N}}$ defined as

$$y_{n+1} = \phi_T(y_n).$$

If we prove that this sequence converges to y^* , then we can deduce that

$$y^* = \phi_T(y^*).$$

This means that there exists a fixed-point for the mapping ϕ_T , which implies in particular that (1) is well-posed in the desired functional spaces. To prove the convergence of this sequence, we prove that it is a Cauchy sequence. Indeed,

since $\mathbf{B}_K(y_0)$ is complete, any Cauchy sequence converges in this set. By induction, one can prove that

$$\|y_{n+1} - y_n\|_T \leq (C(y_0, K)T)^n \|y_0 - y_1\|_T \quad (27)$$

and

$$y_n \in \mathbf{B}_K(y_0). \quad (28)$$

Indeed, thanks to the choice of T in (26), these two properties are easily proved for $n = 0$ and, moreover, we have with (24), for all $n \geq 1$

$$\begin{aligned} \|y_{n+1} - y_n\|_T &= \|\phi(y_n) - \phi(y_{n-1})\|_T \\ &\leq C(y_0, K)T \|\phi(y_{n-1}) - \phi(y_{n-2})\|_T \\ &\leq (C(y_0, K)T)^n \|y_0 - y_1\|_T. \end{aligned} \quad (29)$$

The inequality (27) can be deduced from the above inequality. The property (28) can be proved as follows:

$$\begin{aligned} \|y_{n+1}\| &\leq \|y_n\|_T + \|y_1\|_T + \|y_0\|_T \\ &\leq \|y_0\|_T + \|\phi_T(y_0)\|_T + \|\phi_T(y_0)\|_T + \|y_0\|_T \\ &\leq K \end{aligned} \quad (30)$$

where in the first line we have used (27) and, in the second line, we have used the fact that $y_n \in \mathbf{B}_K(y_0)$.

The two properties (27) and (28) show that the sequence y_n is a Cauchy sequence. Since $\mathbf{B}_K(y_0)$ is a complete set, this sequence is therefore convergent. This means in particular that there exists a fixed-point to the mapping $\phi_T(y_0)$, which implies that, for sufficiently small time T , there exists a unique solution $z \in L^\infty(0, T; W^{1,\infty}(0, 1))$ and $z_t \in L^\infty(0, T; L^\infty(0, 1))$. Thanks to (20), we can deduce that there exists unique solution $z \in L^\infty(\mathbb{R}_+; W^{1,\infty}(0, 1))$ and $z_t \in L^\infty(\mathbb{R}_+; L^\infty(0, 1))$. This concludes the proof of Theorem 1. \bullet

3.2 Proof of Theorem 2

The proof of Theorem 2 is divided into two steps: first, we transform the system (1) into a system in the form (A.1), which is in particular a linear time-variant system, and apply Theorem 3 for the space H_2 , which is indeed the only Hilbert space among all the spaces $H_p(0, 1)$. Second, using the fact that the solutions are bounded in $H_\infty(0, 1)$ thanks to Theorem 1, and invoking an interpolation theorem, which is the Riesz-Thorin theorem, we conclude.

Proof.

First step: semi-global exponential stability in H_2 .

We first fix $p = 2$, but still consider the initial conditions $(z_0, z_1) \in H_\infty(0, 1)$. Moreover, we suppose that there exists a positive constant R such that the initial conditions satisfy

$$\|(z_0, z_1)\|_{H_\infty(0,1)} \leq R. \quad (31)$$

In particular, due to Theorem 1, one has

$$\|(z, z_t)\|_{H_\infty(0,1)} \leq 2R, \quad \forall t \geq 0. \quad (32)$$

The system (1) may be rewritten as follows

$$\begin{cases} z_{tt} = z_{xx} - d(t, x)a(x)z_t, \\ z(t, 0) = z(t, 1) = 0, \\ z(0, x) = z_0(x), z_t(0, x) = z_1(x), \end{cases} \quad (33)$$

with

$$d(t, x) = \begin{cases} \frac{\sigma(\sqrt{a(x)}z_t)}{\sqrt{a(x)}z_t}, & \text{if } a(x)z_t \neq 0, \\ C_1, & \text{if } a(x)z_t = 0. \end{cases} \quad (34)$$

This function comes from the function $t \mapsto \frac{\sigma(\sqrt{a(x)}z_t)}{\sqrt{a(x)}z_t}$, extended at 0, which is possible due to the differentiability of the function σ at 0.

This system is in the form (A.1), with $H = H_2$, $U = H_2$, $D(A) = D_2$, and the operators A and B defined as follows

$$\begin{aligned} A : D(A) \subset H &\rightarrow H \\ [v_1 \ v_2]^\top &\mapsto [v_2 \ v_1'']^\top \end{aligned} \quad (35)$$

and $B = [0 \ \sqrt{a(x)}]^\top$. It is well known that A generates a strongly continuous semigroup of contractions (see Haraux (2018)). Now, let us check whether $d(t, \cdot)$ satisfies (A.2), for all $t \geq 0$.

Since the initial conditions $(z_0, z_1) \in H_\infty(0, 1)$ and are bounded by R in the $H_\infty(0, 1)$ norm, then invoking Theorem 1 we have, for all $t \geq 0$

$$\sup_{x \in [0, 1]} |z_t(t, x)| \leq 2R. \quad (36)$$

Moreover, since $\sqrt{a(x)} \leq \sqrt{a_\infty}$, for all $x \in [0, 1]$, and because d is a continuous function, there exist two positive constants d_0 and d_1 , depending on R and a_∞ such that

$$\begin{aligned} d_0 &:= \min_{\xi \in [-2\sqrt{a_\infty}R, 2\sqrt{a_\infty}R]} \frac{\sigma(\xi)}{\xi} \leq d(t, x) \\ &\leq \max_{\xi \in [-2\sqrt{a_\infty}R, 2\sqrt{a_\infty}R]} \frac{\sigma(\xi)}{\xi} := d_1. \end{aligned} \quad (37)$$

Note moreover that the origin of the following system

$$\begin{cases} z_t = z_{xx} - d_0 z_t \\ z(t, 0) = z(t, 1) = 0 \\ z(0, x) = z_0(x), \quad z_t(0, x) = z_1(x). \end{cases} \quad (38)$$

is exponentially stable in H_2 for any initial conditions $(z_0, z_1) \in H_2$ (see e.g. Prieur et al. (2016)). The related operator of this system is $A - d_0 B B^*$, with domain $D(A_{d_0}) = D_2$. Therefore, all the properties required in Theorem 3 are satisfied. Hence, there exist two positive constants $K := K(R)$ and $\beta := \beta(R)$ such that

$$\|(z, z_t)\|_{H_2} \leq K e^{-\beta t} \|(z_0, z_1)\|_{H_2}, \quad \forall t \geq 0. \quad (39)$$

Second step: Semi-global exponential stability in $H_p(0, 1)$. From Theorem 1, we know that, for every initial conditions $(z_0, z_1) \in H_\infty(0, 1)$ satisfying $\|(z_0, z_1)\|_{H_\infty(0, 1)} \leq R$, and noticing that the trajectory of (1) can be expressed with the evolution family $W(t, 0)$ associated to (33)³, one has

$$\|W(t, 0)(z_0, z_1)\|_{H_\infty(0, 1)} \leq 2R, \quad \forall t \geq 0. \quad (40)$$

Now, fix $t > 0$. Note that $W(t, 0)$ is a linear operator from $(L^2(0, 1))^2$ (resp. $(L^\infty(0, 1))^2$) to $(L^2(0, 1))^2$ (resp. $(L^\infty(0, 1))^2$), if it associates $(z'_0, z_1) \in L^2(0, 1)^2$ (resp. $(z'_0, z_1) \in L^\infty(0, 1)^2$) to $(z_x, z_t) \in L^2(0, 1)$ (resp. $(z_x, z_t) \in L^\infty(0, 1)^2$). Hence, we can apply the so-called Riesz-Thorin theorem (Bergh and L ofstr om, 2012, Theorem 1.1.1, Page 2) and conclude that

$$\|W(t, 0)\|_{H_p(0, 1)} \leq 2 \left(\frac{K}{2}\right)^{\frac{2}{p}} e^{-\frac{2\beta}{p}t}, \quad \forall t \geq 0, \quad (41)$$

³ Evolution families are extension of semigroups for time-variant linear infinite-dimensional systems.

where $W(t, 0)$ corresponds exactly to the trajectory of (1) with the initial condition $z_0, z_1 \in H_\infty(0, 1)$. In particular, for every $(z_0, z_1) \in H_\infty(0, 1)$, one therefore has, for every $t \geq 0$

$$\begin{aligned} \|W(t, 0)(z_0, z_1)\|_{H_p(0, 1)} &\leq \|W(t, 0)\|_{H_p(0, 1)} \|(z_0, z_1)\|_{H_p(0, 1)} \\ &\leq 2 \left(\frac{K}{2}\right)^{\frac{2}{p}} e^{-\frac{2\beta}{p}t} \|(z_0, z_1)\|_{H_p(0, 1)}. \end{aligned}$$

This concludes the proof of Theorem 2. •

4. CONCLUSION

In this paper, we have provided a well-posedness analysis of a one-dimensional wave equation subject to a nonlinear nonmonotone damping. Furthermore, a characterization of the asymptotic behavior of the latter system is given. It is proved with Lyapunov techniques. This work paves the way to many others. For instance, a first open and natural question would be the case of multidimensional wave equations, for which there does not exist any proof of well-posedness in the functional setting introduced in this paper.

Appendix A. EXPONENTIAL CONVERGENCE RESULT FOR A LINEAR TIME-VARIANT SYSTEM

This appendix is devoted to the statement and the proof of a theorem dealing with a time-variant linear infinite-dimensional system. Indeed, as it is illustrated in Section 3.2, we can transform (1) as a time-variant linear infinite-dimensional system. This transformation is inspired by Chitour et al. (1995) and Liu et al. (1996), where such a technique is applied for finite-dimensional linear systems subject to a saturation. To define it, let us introduce H (resp. U), a Hilbert space which is equipped with the norm $\|\cdot\|_H$ and the scalar product $\langle \cdot, \cdot \rangle_H$. The system under study in this section is the following:

$$\begin{cases} \frac{d}{dt}v = (A - d(t)BB^*)v := A_d(t)v \\ v(\tau) = v_\tau, \end{cases} \quad (A.1)$$

where $A : D(A) \subset H \rightarrow H$, with $D(A)$ the domain of the operator A that we suppose densely defined in H , $B \in \mathcal{L}(U, H)$, B^* denotes the adjoint of B and d is a continuous function. We assume moreover that A and its adjoint A^* are dissipative, and that there exist positive constants d_0 and d_1 such that d satisfies, for all $t \geq 0$.

$$d_0 \leq d(t) \leq d_1. \quad (A.2)$$

Due to these properties, the domain of $A_d(t)$ is equal to $D(A)$ and, therefore, does not depend on t . Moreover, always because of these properties, and since A and A^* are dissipative, one can prove that, for all $t \geq 0$ and for all $v \in D(A)$

$$\langle A_d(t)v, v \rangle_H \leq 0, \quad \langle A_d^*(t)v, v \rangle_H \leq 0, \quad (A.3)$$

which implies that each operator $A_d(t)$ generates a strongly continuous semigroup of contractions. Hence, due to the discussion provided in (Chicone and Latushkin, 1999, Page 123), there exists an evolution family $(W(\theta, \tau))_{\theta \geq \tau}$ solving (A.1). In particular, this means that there exists a unique strong solution $v \in C(\mathbb{R}_+, D(A))$ to (A.1) for every $v_0 \in D(A)$.

We are now in position to state the following result.

Theorem 3. Consider the system given by (A.1). Supposing that the origin of the following system

$$\begin{cases} \frac{d}{dt}v = (A - d_0BB^*)v := A_{d_0}v, \\ v(0) = v_0 \end{cases} \quad (\text{A.4})$$

is globally exponentially stable, then, for any initial conditions $v_0 \in H$, the origin of (A.1) converges exponentially to 0 with $\tau = 0$.

Remark 1. The global asymptotic stability is not ensured for the linear time-variant system (A.1). Indeed, for such systems, the global asymptotic stability has to be proven for every initial condition v_τ and for every $\tau \geq 0$. This is not an issue for us. Indeed, we use this result to prove the attractivity of (1). The stability of (1) already holds thanks to Theorem 1. \circ

Proof. Since the origin of (A.4) is globally exponentially stable, then, due to Datko (1970), there exist a self-adjoint operator $P \in \mathcal{L}(H)$ and a positive constant C such that the following inequality holds true

$$\langle PA_{d_0}v, v \rangle_H + \langle Pv, A_{d_0}v \rangle_H \leq -C\|v\|_H^2, \quad \forall v \in D(A) \quad (\text{A.5})$$

Moreover, note that A_{d_0} is also a dissipative operator, which means in particular that

$$\langle A_{d_0}v, v \rangle_H + \langle v, A_{d_0}v \rangle_H \leq 0, \quad \forall v \in D(A). \quad (\text{A.6})$$

Now, consider the following candidate Lyapunov functional for (A.1):

$$V(v) := \langle Pv, v \rangle_H + M\|v\|_H^2, \quad (\text{A.7})$$

where M is a positive constant which has to be defined. The time derivative of V along the trajectories of (A.1) yields

$$\begin{aligned} \frac{d}{dt}V(v) &= \langle (P + MI_H)A_{d(t)}v, v \rangle_H + \langle (P + MI_H)v, A_{d(t)}v \rangle_H \\ &= \langle PA_{d_0}v, v \rangle_H + \langle Pv, A_{d_0}v \rangle_H \\ &\quad + M(\langle A_{d_0}v, v \rangle_H + \langle v, A_{d_0}v \rangle_H) \\ &\quad - \langle B(d(t) - d_0I_H)B^*v, (P + MI_H)v \rangle_H \\ &\quad - \langle (P + MI_H)v, B(d(t) - d_0I_H)B^*v \rangle_H. \\ &\leq -C\|v\|_H^2 - d_0M\|B\|_{\mathcal{L}(H,U)}^2\|v\|_H^2 \\ &\quad + 2(d_1 - d_0)\|B^*\|_{\mathcal{L}(H,U)}\|P\|_{\mathcal{L}(H)}\|v\|_H^2 \\ &\quad - 2M\langle (d(t) - d_0)B^*v, B^*v \rangle_U, \end{aligned} \quad (\text{A.8})$$

where we have used (A.2) and (A.6) to get the inequality. Recalling that $d(t) - d_0 \geq 0$, and setting

$$M = \frac{2(d_1 - d_0)\|P\|_{\mathcal{L}(H)}}{d_0\|B\|_{\mathcal{L}(H,U)}},$$

one has

$$\frac{d}{dt}V(v) \leq -C\|v\|_H^2, \quad \forall v \in D(A). \quad (\text{A.9})$$

Since V satisfies the following inequalities

$$M\|v\|_H^2 \leq V(v) \leq (\|P\|_{\mathcal{L}(H)} + M)\|v\|_H^2 \quad (\text{A.10})$$

one can conclude that

$$\|v\|_H^2 \leq \frac{\|P\|_{\mathcal{L}(H)} + M}{M} \exp\left(-\frac{C}{\|P\|_{\mathcal{L}(H)} + M}t\right) \|v_0\|_H^2, \quad (\text{A.11})$$

which ends the proof of Theorem 3. \bullet

REFERENCES

- Alabau-Boussouira, F. (2012). On some recent advances on stabilization for hyperbolic equations. In *Control of partial differential equations*, 1–100. Springer.
- Amadori, D., Aqel, F.A.Z., and Santo, E.D. (2019). Decay of approximate solutions for the damped semilinear wave equation on a bounded 1d domain. URL <https://arxiv.org/abs/1807.01968>.
- Bergh, J. and Löfström, J. (2012). *Interpolation spaces: an introduction*, volume 223. Springer Science & Business Media.
- Chicone, C. and Latushkin, Y. (1999). *Evolution semigroups in dynamical systems and differential equations*. 70. American Mathematical Soc.
- Chitour, Y., Liu, W., and Sontag, E. (1995). On the continuity and incremental-gain properties of certain saturated linear feedback loops. *International Journal of Robust and Nonlinear Control*, 5(5), 413–440.
- Datko, R. (1970). Extending a theorem of A.M. Liapunov to Hilbert space. *Journal of Mathematical analysis and applications*, 32(3), 610–616.
- Feireisl, E. (1993). Strong decay for wave equations with nonlinear nonmonotone damping. *Nonlinear Analysis: Theory, Methods & Applications*, 21(1), 49–63.
- Haraux, A. (2009). L^p estimates of solutions to some nonlinear wave equations in one space dimension. *International Journal of Mathematical Modelling and Numerical Optimisation*, 1(1-2), 146–152.
- Haraux, A. (2018). *Nonlinear vibrations and the wave equation*. SpringerBriefs in Mathematics.
- Kang, W. and Fridman, E. (2017). Boundary control of delayed ode–heat cascade under actuator saturation. *Automatica*, 83, 252–261.
- Liu, W., Chitour, Y., and Sontag, E. (1996). On finite-gain stabilizability of linear systems subject to input saturation. *SIAM Journal on Control and Optimization*, 34(4), 1190–1219.
- Martinez, P. and Vancostenoble, J. (2000). Exponential stability for the wave equation with weak nonmonotone damping. *Portugaliae Mathematica*, 57(3), 285–310.
- Marx, S., Andrieu, V., and Prieur, C. (2017a). Cone-bounded feedback laws for m -dissipative operators on Hilbert spaces. *Mathematics of Control, Signals and Systems*, 29(4), 18.
- Marx, S., Cerpa, E., Prieur, C., and Andrieu, V. (2017b). Global stabilization of a Korteweg-de Vries equation with a saturating distributed control. *SIAM Journal on Control and Optimization*, 55(3), 1452–1480.
- Marx, S., Chitour, Y., and Prieur, C. (2018). Stability analysis of dissipative systems subject to nonlinear damping via Lyapunov techniques. *arXiv preprint arXiv:1808.05370*.
- Prieur, C., Tarbouriech, S., and Gomes da Silva Jr, J.M. (2016). Wave equation with cone-bounded control laws. *IEEE Trans. on Automat. Control*, 61(11), 3452–3463.
- Slemrod, M. (1989). Feedback stabilization of a linear control system in Hilbert space with an a priori bounded control. *Mathematics of Control, Signals and Systems*, 2(3), 847–857.
- Strauss, W.A. (1992). *Partial differential equations*. John Wiley & Sons New York, NY, USA.